

2nd-Order ODEs & Equilibriums

Objectives:

1. Introducing linear 2nd-order ODEs
2. Verifying solutions to linear 2nd-order ODEs
3. Equilibrium solutions to linear 2nd-order ODEs

Recall: Exponential Growth/Decay

$$\frac{dy}{dt} = ry \quad \leftarrow \text{1st-order, linear, homo, auto}$$

r is a parameter
separation of variables

$$\int \frac{dy}{y} = \int r dt$$


$$\ln(y) = rt + C$$

$$y(t) = e^{rt+C}$$

$$y(t) = Ae^{rt}$$

$$\text{Verify: } \left. \begin{array}{l} y = Ae^{rt} \\ y' = A r e^{rt} \end{array} \right\} \frac{dy}{dt} = ry$$
$$\cancel{A} e^{rt} = r \cancel{A} e^{rt}$$
$$1 = 1$$

Falling Object



Newton's 2nd Law of Motion: $\vec{F} = m\vec{a}$

- $\vec{F}$ is the net force
- m is the mass of the object
- a is the acceleration

Assumptions:

- No air resistance

- gravity is the only force acting on the object (vertical fall)
- the weight is $\vec{F}$ because gravity is the only force and direction is downwards
- so, $\cancel{m}g = \cancel{m}a$
 $g = a \rightarrow \text{gravity} = \text{acceleration regardless of mass}$
- Object is near earth surface

→ Let $y(t)$ be the height of the object,
 $y'(t)$ be the velocity, and
 $y''(t)$ be the acceleration.

Since y'' is the acceleration and $a = g$, then $\vec{F} = \vec{F}_{\text{gravity}}$ or $y'' = g$.

→ the 2nd-order ODE for modeling falling object is

$$y'' = g \leftarrow \text{2nd-order, linear, non-homo, auto}$$

→ Solution using direct integration: $\int y'' = \int g dt$

$$y' = gt + C_1 \leftarrow \text{velocity}$$

$$\int y' = \int (gt + C_1) dt$$

$$y(t) = \frac{gt^2}{2} + \underbrace{C_1 t + C_2}_{\text{height}} \leftarrow \text{height}$$

two constants because we integrated twice.

→ You need two initial conditions $y(0) = y_0 \leftarrow \text{initial height}$
 $y'(0) = 0 \leftarrow \text{initial velocity (held up at } y_0 \text{ height)}$

→ Solution graph:


gravity is $g < 0$
 (negative)
 Earth $g = -9.8 \text{ m/s}^2$



$$t_{\text{floor}} = \frac{C_1}{g} + \frac{\sqrt{2C_2 g + C_1^2}}{g} \quad \text{for } 0 = \frac{gt^2}{2} + C_1 t + C_2$$

time it takes to reach floor

Falling Object with Air Resistance



With the same assumptions at the Falling object example.

- Additional assumptions:
- With air resistance (drag)
 - Air resistance $\vec{F}_{\text{resist}}$ is the force acting against gravity which is proportional to the velocity.
 - $\vec{F}_{\text{resist}} = -ky' \leftarrow \text{for low velocity}$
or
 $\vec{F}_{\text{resist}} = -k(y')^2 \leftarrow \text{for high velocities}$
 $\hookrightarrow k$ is the resistant coefficient

The total force is still $\vec{F} = m\vec{a}$ or $\vec{F} = m y''$.

→ ODE: $\vec{F} = \vec{F}_{\text{gravity}} + \vec{F}_{\text{resist}}$

$$\downarrow$$
$$m y'' = g - k y'$$

or

$$m y'' + k y' = g \quad \leftarrow \text{2nd-order, linear, auto, non-homo}$$

↓

We can't solve this using direct integration but we can using other methods

Verifying Solutions to 2nd-order Linear ODEs

* Example 1: $y'' + 2y = 0 \rightarrow$ proposed solutions $y_1 = \sin(\sqrt{2}t)$
 $y_2 = \cos(\sqrt{2}t)$

• Verify $y_1 = \sin(\sqrt{2}t)$
 $y_1' = \sqrt{2} \cos(\sqrt{2}t)$
 $y_1'' = -2 \sin(\sqrt{2}t)$

$$y'' + 2y = 0 \rightarrow -2 \sin(\sqrt{2}t) + 2 \sin(\sqrt{2}t) = 0$$
$$0 = 0 \quad \checkmark \quad \text{So, } y_1 \text{ is a solution.}$$

• Verify $y_2 = \cos(\sqrt{2}t)$
 $y_2' = -\sqrt{2} \sin(\sqrt{2}t)$
 $y_2'' = -2 \cos(\sqrt{2}t)$

$$y'' + 2y = 0 \rightarrow -2 \cos(\sqrt{2}t) + 2 \cos(\sqrt{2}t) = 0$$
$$0 = 0 \quad \text{So, } y_2 \text{ is a solution}$$

* Example 2: $y'' - y = 0 \rightarrow$ proposed solutions $y_1 = e^t$
 $y_2 = e^{-t}$

• Verify $y_1 = e^t$
 $y_1' = e^t$
 $y_1'' = e^t$

$$y'' - y = 0 \rightarrow e^t - e^t = 0$$
$$0 = 0 \quad \text{So, } y_1 \text{ is a solution.}$$

• Verify $y_2 = e^{-t}$
 $y_2' = -e^{-t}$
 $y_2'' = e^{-t}$

$$y'' - y = 0 \rightarrow e^{-t} - e^{-t} = 0$$
$$0 = 0 \quad \text{So, } y_2 \text{ is a solution}$$

$$y'' - y = 0 \rightarrow \cancel{e^t} - \cancel{e^{-t}} = 0$$

$$0 = 0 \text{ so, } y_2 \text{ is a solution.}$$

* Example 3: $y'' + 3y' = 0 \rightarrow$ proposed solution $y = e^{-3t}$

• verify

$$y = e^{-3t}$$

$$y' = -3e^{-3t}$$

$$y'' = 9e^{-3t}$$

$$y'' + 3y' = 0 \rightarrow 9e^{-3t} + 3(-3e^{-3t}) = 0$$

$$\cancel{9e^{-3t}} - \cancel{9e^{-3t}} = 0$$

$$0 = 0$$

Equilibrium Solutions of 2nd-order Linear ODEs

An equilibrium solution is a constant solution y where $y' = 0$ and $y'' = 0$.

* Example 4: $y'' - y = 0 \rightarrow$ set $y'' = 0$.

$$0 - y = 0 \rightarrow y = 0 \text{ a constant}$$

So, an equilibrium solution.

* Example 5: $y'' + 3y' = 0 \rightarrow$ set $y' = 0$ & $y'' = 0$.

$$0 + 3(0) = 0 \rightarrow 0 = 0, \text{ meaning an equilibrium exists}$$

but we can't know it... yet.